

Pairing symmetry transitions in the even-denominator FQHE system

Kentaro Nomura* and Daijiro Yoshioka

Department of Basic Science, University of Tokyo, 3-8-1 Komaba, Tokyo 153-8902

(February 1, 2008)

Transitions from a paired quantum Hall state to another quantum Hall state in bilayer systems are discussed in the framework of the edge theory. Starting from the edge theory for the Haldane-Rezayi state, it is shown that the charging effect of a bilayer system which breaks the $SU(2)$ symmetry of the pseudo-spin shifts the central charge and the conformal dimensions of the fermionic fields which describe the pseudo-spin sector in the edge theory. This corresponds to the transition from Haldane-Rezayi state to Halperin's 331 state, or singlet d-wave to triplet p-wave ABM type paired state in the composite fermion picture. Considering interlayer tunneling, the tunneling rate-capacitance phase diagram for the $\nu = 5/2$ paired bilayer system is discussed.

A bilayer quantum Hall system can be mapped to an equivalent spin-1/2 system by assigning $\uparrow(\downarrow)$ pseudo spins to electrons in the upper (lower) layer, [1] where the actual electron spins are assumed to be polarized. At finite layer separation, the Hartree part of the electron-electron repulsive interaction produces a local capacitive charging energy that is minimized when the two layers have equal electron density. Especially for the $\nu = 1/2$ (odd integer) quantum Hall ferromagnet, the expectation value of the z-component of the pseudo spin vanishes and the pseudo spin system has an easy-plane anisotropy that gives the itinerant ferromagnet with an XY symmetry in the absence of interlayer tunneling. The characteristic parameters are the distance between the layers, the interlayer tunneling rate, and the electric capacitance etc. The grand state is characterized by these parameters. The simplest Abelian quantum Hall state for the bilayer system is described by the two-component generalization of the Laughlin wave function first introduced by Halperin as [2]

$$\Psi_{mmn} = \prod_{i < j} (z_i^\uparrow - z_j^\uparrow)^m \prod_{i < j} (z_i^\downarrow - z_j^\downarrow)^m \prod_{i, j} (z_i^\uparrow - z_j^\downarrow)^n, \quad (1)$$

where we abbreviated the factor $e^{-\frac{1}{4} \sum (|z_i^\uparrow|^2 + |z_i^\downarrow|^2)}$. In the case of $\nu = 1/2$, $(m, m, n) = (3, 3, 1)$, and the wave function is rewritten as

$$\Psi_{331} = \det \left(\frac{1}{z_i^\uparrow - z_j^\downarrow} \right) \Psi_{222}. \quad (2)$$

This state is called 331-state.

Haldane and Rezayi have proposed a two-component singlet wave function to explain the quantum Hall effect at $\nu = 5/2$. [3,4] This state is an exact grand state of the hollow-core model and is given by

$$\Psi_{\text{HR}} = \det \left(\frac{1}{(z_i^\uparrow - z_j^\downarrow)^2} \right) \Psi_{222}, \quad (3)$$

which is called Haldane-Rezayi(HR) state.

In above two states the electron spins are unpolarized. Moore and Read proposed spin polarized state which is described by the trial function; [5]

$$\Psi_{\text{Pf}} = \text{Pf} \left(\frac{1}{z_i - z_j} \right) \prod_{i < j} (z_i - z_j)^2, \quad (4)$$

This state is called Pfaffian state. $\text{Pf}(M_{ij})$ means $A \prod_i M_{2i-1, 2i}$. Greiter et al. studied the Pfaffian state as a paired state of the composite fermions. [6,7] Ho considered interesting connections between the internal order in these types of wave functions and analogous order in superfluid ^3He . [8] The order parameter symmetry is triplet p-wave with ABM and A_1 type in ^3He for 331 and Pfaffian state, respectively. Similarly HR state can be regarded as d-wave paired state. [9]

The wave functions for the grand state and the excited states in these paired states are interpreted as the correlation functions of proper conformal field theories(CFT). In the case of the Pfaffian, 331, and HR states, corresponding theories have a central charge $c = 1 + 1/2, 1 + 1, 1 - 2$. [5,10-12] These conformal theories also describe the edge excitations on the boundaries of the sample. In the even-denominator quantum Hall regime such as $\nu = 1/2$ and $5/2$, the grand state was used to be considered as one of these three states. Recently, Read and Green argued that the HR state is right at bulk transition point between weak and strong pairing d-wave states. [9] In this paper we discussed another direction of the developments which is the grand state of the bilayer systems. Let us consider the charging effect and interlayer tunneling which are described in terms of the pseudo spin as

$$H_U = \epsilon_c \int d^2 \mathbf{x} (S^z(\mathbf{x}))^2, \quad (5)$$

and

$$H_T = -\Delta_{SAS} \int d^2 \mathbf{x} S^x(\mathbf{x}), \quad (6)$$

respectively. Latter is regarded as Zeeman term for pseudo-spin. It is well known that introducing a Zeeman term the pseudo-spins align to the direction of the (pseudo) magnetic field, and pseudo-spin polarization transition occurs. This process corresponds to the 331-Pfaffian transition which is argued in Ref 8). On the other hand, we can expect that if the introduced anisotropy is XY (i.e. easy-plane) type, the singlet-triplet transition might appear.

In this paper we describe the latter transition which occurs between the HR and 331 due to the charging effect in the bilayer system with hollow-core interaction which is perturbed by layer separation in the framework of the edge theory. Our results predict that the 331 state is favored for the charging term. This is consistent with a numerical studies [13,14] for realistic interaction.

The edge theory is first established by X.G. Wen [15,16] and the connection with the bulk wave function is indicated by Moore and Read. [5]

First we review this connection for the $\nu = 1/q$ Laughlin state. The grand state wave function is written as the correlation function of the $c = 1$ CFT.

$$\Psi_{\text{Ln}} = \langle \prod_{i=1}^{N_e} \psi_e(z_i) \Phi_{\Delta} \rangle, \quad (7)$$

where $\psi_e(z) =: e^{i\sqrt{q}\phi(z)} :$ is the electron field operator, $\Phi_{\Delta} =: e^{-i\rho_0 \int d^2z \sqrt{q}\phi(z)} :$ is the primary field which has the conformal dimension $\Delta = qN_e/2$, and we ignore the singular phase which is derived from the right hand side of eq.(7). The $c = 1$ CFT is represented by the action of the free bosonic field

$$S = \int d^2x \frac{1}{2\pi} \partial\phi \bar{\partial}\phi, \quad (8)$$

where z is the complex coordinate, the correlation exponent of the theory is given as $K = \nu$. The edge excitations are connected with the angular momentum excitations of the bulk. For a disc geometry, the Hamiltonian for edge modes is given as

$$H = \frac{2\pi v}{L} L_0, \quad (9)$$

where $L_0 = \frac{1}{2}J_0^2 + \sum_{n>1} J_n J_{-n}$ is the holomorphic part of the zero-component of the Virasoro algebra which can be regarded as the z-component of the total orbital angular momentum, J is the Kac-Moody current, and we ignore the correction term which is proportional to the central charge.

Now we come back to the present bilayer case. The edge theory of the paired quantum Hall state has an internal degree of freedom which corresponds to the pseudo spin in the bilayer systems. The additional prefactor to Laughlin-Jastrow type factor of the three paired state wave functions in eqs.(2)-(4) arise from these pseudo spin mode.

The edge theory for the 331 state is described by the Hamiltonian

$$H_{331} = \frac{1}{4\pi} \int dx v_{IJ} : \partial_x \phi_I \partial_x \phi_J :, \quad (10)$$

where v_{IJ} is a symmetric matrix which depends on the confining potential and interchannel interactions at the edge. [5,16] The commutation relations of the bosonic fields are $[\phi_I(x), \phi_J(x')] = i\pi K_{IJ} \text{sgn}(x - x')$, where K is the K-matrix given as [17]

$$K = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix}. \quad (11)$$

In the charge-pseudo spin basis, $\phi_c = \frac{1}{\sqrt{v}}(\phi_{\uparrow} + \phi_{\downarrow})$, $\phi_s = \phi_{\uparrow} - \phi_{\downarrow}$, the two components can be separated as

$$H_{331} = \frac{1}{4\pi} \int dx \left[v_c : (\partial_x \phi_c)^2 : + v_s : (\partial_x \phi_s)^2 : \right]. \quad (12)$$

Both charge and spin sectors are also written as the free bosonic form which have the correlation exponent $K_c = \nu, K_s = 1$, respectively. The electron operator is given by $\psi_{e\uparrow} = e^{\frac{1}{\sqrt{v}}\phi_c + i\phi_s}$, $\psi_{e\downarrow} = e^{\frac{1}{\sqrt{v}}\phi_c - i\phi_s}$. Because the case $K_s = 1$ is equivalent to the free fermion theory, the spin sector Hamiltonian is also described by

$$H_{331}^s = -iv \int dx \psi^\dagger \partial_x \psi = \frac{2\pi v}{L} L_0, \quad (13)$$

where the fermionic field ψ is related to the bosonic field by $\psi = e^{i\phi_s}$, and $L_0 = \frac{1}{2}(J_0^z)^2 + \sum_{n>1} J_n^z J_{-n}^z$, $J^z = i\partial\phi_s$ is the z-component of the pseudo-spin current density on the edge.

As we mentioned above, the spin sector corresponds to the degrees of the freedom of the pairing. Actually the equality

$$\det \left(\frac{1}{z_i^\uparrow - z_j^\downarrow} \right) = \langle \prod_{i,j} \psi^\dagger(z_i^\uparrow) \psi(z_j^\downarrow) \rangle \quad (14)$$

can be derived.

In the HR state, the electron operator is written as $\psi_{e\sigma} = \partial\theta_\sigma e^{i\sqrt{2}\phi_c}$, The determinant part of the wave function is reconstructed by the conformal field theory of the fermionic ghost field as

$$\det \left(\frac{1}{(z_i^\uparrow - z_j^\downarrow)^2} \right) = \langle \prod_{i,j} \partial\theta_\uparrow(z_i^\uparrow) \partial\theta_\downarrow(z_j^\downarrow) \rangle,$$

that corresponds to the spin sector of the edge excitation. On the other hand, the charge sector is same as the theory for the 331 state. The spin sector of the edge excitations is described by

$$\begin{aligned} \theta_\uparrow &= \sqrt{\frac{2\pi}{L}} \sum_{k>0} \frac{1}{\sqrt{k}} (c_{k\uparrow} e^{ikx} + c_{k\downarrow} e^{-ikx}) \\ \theta_\downarrow &= \sqrt{\frac{2\pi}{L}} \sum_{k>0} \frac{1}{\sqrt{k}} (c_{k\uparrow} e^{ikx} - c_{k\downarrow} e^{-ikx}). \end{aligned} \quad (15)$$

The Hamiltonian is given as [11,12]

$$H_{\text{HR}}^s = \int dx \frac{v_s}{4\pi} : (\partial_x \theta_{\uparrow} \partial_x \theta_{\downarrow} + \partial_x \theta_{\uparrow}^{\dagger} \partial_x \theta_{\downarrow}^{\dagger}) : \\ = \sum_{k>0} v_s k \left(c_{k\uparrow}^{\dagger} c_{k\uparrow} + c_{k\downarrow}^{\dagger} c_{k\downarrow} \right), \quad (16)$$

which has the global SU(2) symmetry with the total spin generator

$$S_z = \frac{1}{2} \sum_{k>0} (c_{k\uparrow}^{\dagger} c_{k\uparrow} - c_{k\downarrow}^{\dagger} c_{k\downarrow}), \\ S_+ = \sum_{k>0} c_{k\uparrow}^{\dagger} c_{k\downarrow}, \quad S_- = \sum_{k>0} c_{k\downarrow}^{\dagger} c_{k\uparrow}. \quad (17)$$

To go on our argument further, we convert to the (ξ, η) -representation:

$$S_{\text{HR}}^s = \int d^2x \frac{1}{2\pi} (\eta \bar{\partial} \xi + \bar{\eta} \partial \bar{\xi}), \quad (18)$$

where the field $\eta = \partial \theta_{\uparrow}$ and $\xi = \theta_{\downarrow}$ are fermionic field which satisfy

$$\langle \xi(z) \eta(\omega) \rangle = \frac{1}{z - \omega}. \quad (19)$$

The conformal dimensions of the ξ and η are 0, 1, respectively. The energy-momentum tensor is derived from (18) as

$$T(z) = : \partial \xi \eta(z) : \\ = -\frac{1}{2} : (\eta \partial \xi - \partial \eta \xi)(z) : + \frac{1}{2} : (\partial \xi \eta + \xi \partial \eta)(z) : \\ = \frac{1}{2} J^z(z)^2 + \frac{1}{2} \partial J^z(z), \quad (20)$$

where $J^z(z) = : \xi \eta(z) :$. Calculating the operator product expansion;

$$T(\omega) \eta(z) = \frac{1}{(\omega - z)^2} \eta(z) + \frac{\partial \eta(z)}{\omega - z} + \dots, \quad (21)$$

$$T(\omega) \xi(z) = \frac{0}{(\omega - z)^2} \xi(z) + \frac{\partial \xi(z)}{\omega - z} + \dots, \quad (22)$$

$$T(\omega) T(z) = \frac{-1}{(\omega - z)^4} + \frac{2T(z)}{(\omega - z)^2} + \dots, \quad (23)$$

we can check the conformal dimension of the field η and ξ and the central charge $c = -2$ directly. The zero-component of the energy-momentum tensor is

$$L_0^{c=-2} = \frac{1}{2} J_0^z (J_0^z - 1) + \sum_{n>0} J_n^z J_{-n}^z. \quad (24)$$

Then the spin sector of the edge Hamiltonian for HR-state eq.(16) is also written as

$$H_{\text{HR}}^s = \frac{2\pi v}{L} L_0^{c=-2}. \quad (25)$$

Now we consider the charging effect. On the edge, the charging Hamiltonian (5) is rewritten as

$$H_U = \int dx g J^z(x)^2 \quad (26)$$

where a constant g is proportional to ϵ_c in (5).

The total Hamiltonian is given by

$$H = H^c + H_{\text{HR}}^s + H_U \\ = H^c + \frac{2\pi v_s}{L} \left(\frac{1}{2} (J_0^z)^2 + \sum_{n>0} J_n^z J_{-n}^z - \alpha J_0^z \right). \quad (27)$$

where $v_s = v + gL/2\pi$ is the renormalized velocity, and $\alpha = v/(v + gL/2\pi)$. For a large g , the last term of the eq.(27) can be ignored and the spin sector is modified as below. The energy-momentum tensor which derive the Hamiltonian eq.(27) with $\alpha = 0$ is

$$T_{\text{eff}}(z) = \frac{1}{2} (J^z(z))^2 = -\frac{1}{2} : (\eta \partial \xi - \partial \eta \xi) :$$

This mechanism is called spectral flow which originally discussed in Ref [18–22]. The conformal dimensions of the fermion fields become $\Delta_{\xi} = \Delta_{\eta} = 1/2$. Using Wick's theorem we can check it in the operator product expansion:

$$T_{\text{eff}}(\omega) \eta(z) = \frac{1/2}{(\omega - z)^2} \eta(z) + \frac{\partial \eta(z)}{\omega - z} + \dots, \quad (28)$$

$$T_{\text{eff}}(\omega) \xi(z) = \frac{1/2}{(\omega - z)^2} \xi(z) + \frac{\partial \xi(z)}{\omega - z} + \dots. \quad (29)$$

Not only the conformal dimensions of the fermion fields but the central charge of the spin sector is also shifted from $c = -2$ to unity. So we can rewrite the fermionic field as

$$\psi \equiv \eta = \sqrt{\frac{2\pi}{L}} \sum f_k e^{ikx} \\ \psi^{\dagger} \equiv \xi = \sqrt{\frac{2\pi}{L}} \sum f_k^{\dagger} e^{-ikx}, \quad (30)$$

and the Hamiltonian as

$$H_{\text{eff}}^s = \int \frac{dx}{2\pi} \psi^{\dagger} \left(-iv_s \frac{\partial}{\partial x} \right) \psi \\ = \sum_k v_s k f_k^{\dagger} f_k. \quad (31)$$

In eq.(31), the global SU(2) symmetry represented by eq.(17) is reduced to U(1) XY symmetry. This modified theory can be reconciled with the spin sector for the 331 state. Therefore due to the charging effect, the HR-331 transition occurs.

Next we discuss the effect of interlayer tunneling (6) on the edge that causes the pseudo-spin polarization to x

direction. So the Hamiltonian (16) become that for the single component

$$H_{\text{Pf}}^s = \sum_{k>0} v_s k c_{k+}^\dagger c_{k+} = \int \frac{dx}{2\pi} \chi \left(-iv_s \frac{\partial}{\partial x} \right) \chi, \quad (32)$$

where $c_{k\pm} = \frac{1}{\sqrt{2}}(c_{k\uparrow} \pm c_{k\downarrow})$ and

$$\chi = \sqrt{\frac{2\pi}{L}} \sum_{k>0} (c_{k+} e^{ikx} + c_{k+}^\dagger e^{-ikx}). \quad (33)$$

The spin sector χ have the correlation function

$$\text{Pf} \left(\frac{1}{z_i - z_j} \right) = \langle \chi(z_1) \cdots \chi(z_N) \rangle, \quad (34)$$

that contributes to the bulk wave function (4). So these process is regarded as the polarization transition between the HR and Pfaffian state. A edge theory discription of the 331-Pfaffian transition is discussed in Ref. [9].

Finally we consider the tunneling amplitude versus layer separation d phase diagram. We regard these two parameter as independent. When the interlayer tunneling rate is large. the system has effectively one component. (i.e. pseudo spin is polarized to the easy-plane.) This is the Pfaffian state. Contrary, in the regime where the tunneling amplitude is small, the 331 and HR state are expected to appear. According to above analysis, the HR-331 transition occurs as the charging effect is increased. These statements are summarized as Fig.1.

In this paper we have described the pairing symmetry transition in the bilayer even denominator quantum Hall states in the framework of the conformal field theories. Starting from the edge theory of the HR state the charging term shifts the central charge from $c = -2$ to $c = 1$ and the conformal dimensions of the fermion fields. This is interpreted as the HR-331 transition. In view of the composite fermion theory, HR state is the d-wave paired state and 331 state is the p-wave ABM state. So the transition is from singlet to triplet. Our result shows that the 331 state is favored in the bilayer systems with the finete layer separation. In the context of our edge theory description, we cannot know what kind of the transition this belongs. The detailed investigation of the transitional or crossover points is left to future problems.

ACKNOWLEDGEMENTS

We are grateful to A. Yamaguchi and K. Hamada for useful discussions and thier collaborations. We also acknowledge K. Ino for pointing out the mechanism of the

spectral flow and N. Read for helpful indication for our phase diagram. This work is supported by a Grant-in-Aid for Scientific Research (C) 10640301 from the Ministry of Education, Science, Sports, Culture and Technology.




FIG. 1. The proposed phase diagram where d is separation between the layers. The polarized state is the Pfaffian state only. The unpolarized states are 331 and HR state. The latter state exists as a critical point (dot in the diagram).

* nomura@sola.c.u-tokyo.ac.jp

- [1] J. P. Eisenstein in " *Perspectives in Quantum Hall Effects* " ed. by S. Das Sarma and A. Pinczuk, 1997, John Wiley and Sons, New York.
- [2] B. I. Halperin *Helv. Phys. Acta.* **56** (1983) 75.
- [3] F. D. M. Haldane, R. Rezayi: *Phys. Rev. Lett.* **60** (1988) 956.
- [4] D. Yoshioka, A. H. MacDonald, and S. M. Girvin: *Phys. Rev. B* **38** (1988) 3636.
- [5] G. Moore, N. Read: *Nucl. Phys. B* **360** (1991) 362.
- [6] M. Greiter, X. G. Wen, and F. Wilczek: *Nucl. Phys. B* **374** (1992) 567.
- [7] J. K. Jain: *Phys. Rev. Lett.* **63** (1989) 199.
- [8] T. L. Ho: *Phys. Rev. Lett.* **75** (1995) 1186.
- [9] N. Read and D. Green: *Phys. Rev. B* **61** (2000) 10267.
- [10] X.G. Wen and Y. S. Wu: *Nucl. Phys. B* **419** (1994) 455.
- [11] M. Milovanovic and N. Read: *Phys. Rev. B* **53** (1996) 13559.
- [12] J. C. Lee and X.G. Wen: *Nucl. Phys. B* **542** (1999) 647.
- [13] D. Yoshioka, A. H. MacDonald and S. M. Girvin: *Phys. Rev. B* **39** (1989) 1932.
- [14] V.W. Scarola and J.K. Jain: *preprint cond-mat/0011511*.
- [15] X. G. Wen: *Phys. Rev. B* **41** (1990) 12838.
- [16] X. G. Wen: *Int. J. Mod. Phys. B* **6** (1992) 1711.
- [17] X.G. Wen and A. Zee: *Phys. Rev. B* **46** (1992) 2290.
- [18] V. Gurarie, M. Flohr and Nayak: *Nucl. Phys. B* **498** (1997) 513.
- [19] S. Guruswamy and A. W. W. Ludwig: *Nucl. Phys. B* **519** (1998) 661.
- [20] K. Ino: *Nucl. Phys. B* **532** (1998) 782.
- [21] K. Ino: *Phys. Rev. Lett.* **82** (1999) 4903.
- [22] A. Cappelli, L.S. Georgiev, and I. T. Todorov: *Commun. Math. Phys.* **205** (1999) 657.